CLINOVATE
Advances in Medicine and Healthcare
eISSN: 3107-4421

Volume 2, Issue 1, Jan-Apr 2026
DOI: 10.5281/zenodo.19673611

Perspective

Infant Weaning Practices in Rural India: Bridging the Gap Between Knowledge, Culture, and Practice

Saumya Awasthi[1*]

[1]*Medical Officer, Madhav Child Care Centre, Nagpur, MH, India*

Abstract

Infant weaning, or the transition from exclusive breastfeeding to complementary feeding, is a critical determinant of child growth and development. Despite well-established global and national guidelines, suboptimal weaning practices remain highly prevalent in rural India, contributing to undernutrition and adverse health outcomes. Existing interventions have largely focused on improving caregiver awareness; however, the persistence of inappropriate feeding behaviors suggests that knowledge alone is insufficient to drive meaningful change.

This perspective examines the gap between recommended and actual practices, emphasizing the role of socio-cultural, familial, economic, and health system factors in shaping infant feeding behaviors. Cultural beliefs regarding food suitability, intra-household decision-making dynamics, limited maternal autonomy, and structural constraints such as food insecurity and time limitations significantly influence the adoption of recommended practices. Additionally, standardized and context-insensitive health communication further limits the effectiveness of current interventions.

Addressing these challenges requires a shift toward culturally informed and context-sensitive strategies. Engaging family members, strengthening the communication skills of frontline health workers, and promoting community-based behavioral interventions can enhance the acceptability and sustainability of recommended practices. Integrating scientific guidance with local realities is essential to bridging the gap between knowledge and practice and improving infant nutritional outcomes in rural India.

Keywords: *Infant weaning, complementary feeding, rural India, cultural practices, child nutrition*

How to Cite: Awasthi S. Infant Weaning Practices in Rural India: Bridging the Gap Between Knowledge, Culture, and Practice. CLINOVATE: Advances in Medicine and Healthcare 2026 Apr 24; 2(1). DOI: 10.5281/zenodo.19673611

Introduction

Infant weaning, defined as the transition from exclusive breastfeeding to complementary feeding, is a critical determinant of growth, development, and survival, particularly in resource-limited settings.[1] Global and national guidelines, including those by the World Health Organization, recommend the initiation of complementary feeding at six months of age, alongside continued breastfeeding, with an emphasis on dietary diversity, appropriate frequency, and nutritional adequacy.[2] These recommendations are well established and widely disseminated through public health programs.

Despite this, suboptimal weaning practices remain highly prevalent in rural India.[3] Delayed initiation of complementary feeding, early introduction of nutritionally inadequate foods, and insufficient dietary diversity continue to contribute to undernutrition and adverse health outcomes among infants. The persistence of these practices, even in regions where awareness initiatives have been implemented, suggests that the issue extends beyond a simple lack of knowledge.

Most existing interventions have focused predominantly on educating mothers about recommended feeding practices.[4] However, infant feeding behaviors are not shaped by knowledge alone; they are deeply embedded within a complex interplay of cultural beliefs, family dynamics, socioeconomic constraints, and health system interactions. These contextual factors often influence how information is interpreted, negotiated, and ultimately translated into practice.

This perspective argues that improving infant weaning practices in rural India requires moving beyond information-centric approaches toward a more nuanced understanding of the socio-cultural environment in which caregiving decisions are made. Recognizing and integrating these contextual determinants is essential to bridging the persistent gap between knowledge and practice.

Received: 28/01/2026, Accepted: 17/03/2026, Published: 24/04/2026
*Correspondence and requests should be addressed to S.A. via saumyaawasthi1101@gmail.com

Knowledge–Practice Gap: Beyond Awareness

Over the past decade, awareness regarding optimal infant feeding practices has improved across many rural communities in India.[5] Caregivers are often familiar with recommended timelines for initiating complementary feeding. However, this awareness does not consistently translate into practice. The persistence of inappropriate weaning behaviors highlights a critical gap between knowledge and action, suggesting that information alone is insufficient to drive behavioral change.

Caregivers frequently navigate multiple and sometimes conflicting sources of information, including advice from healthcare providers, family members, and traditional community practices. In such contexts, culturally familiar approaches often take precedence over biomedical recommendations, not necessarily due to resistance, but because they are perceived as more practical, trustworthy, or contextually relevant.

Cultural Beliefs and Food Perceptions

Cultural frameworks play a central role in shaping infant feeding practices.[6] Concepts such as the classification of foods as "hot" or "cold," or beliefs regarding digestibility and suitability for infants, strongly influence decisions about the type and timing of food introduction. Nutrient-dense foods such as pulses, eggs, and certain vegetables are often delayed due to concerns about intolerance or illness, while diluted cereals or liquids may be introduced earlier because they are perceived as safer and easier to digest.

While these practices are rooted in long-standing traditions and caregiving logic, they may inadvertently compromise dietary quality during a critical window of growth and development. Importantly, these beliefs are not inherently irrational but reflect culturally constructed understandings of health and nutrition.

Family Dynamics and Decision-Making Authority

Infant feeding decisions in rural households are rarely made by mothers alone.[7,8] Elder family members, particularly grandmothers, often play a dominant role in guiding childcare practices. Their influence, grounded in experience and tradition, can outweigh formal medical advice.

This dynamic may create tension when recommended practices conflict with established norms. Mothers, especially younger or less empowered ones, may have limited autonomy to implement evidence-based feeding strategies. Additionally, gender roles and household responsibilities constrain the time and capacity of mothers to prepare diverse and age-appropriate foods, further shaping feeding practices.

Socioeconomic and Environmental Constraints

Feeding practices are also closely linked to economic realities and environmental conditions.[1,9] Limited access to diverse and affordable food options restricts the ability of families to follow recommended dietary guidelines. Seasonal variability in food availability, financial constraints, and competing household priorities often dictate food choices.

Even when caregivers are aware of appropriate practices, practical barriers such as food insecurity and time constraints may hinder implementation. In such situations, feeding decisions are frequently driven by feasibility and convenience rather than nutritional adequacy, underscoring the structural nature of the problem.

Health System Limitations and Communication Gaps

Frontline healthcare providers serve as a primary source of information on infant feeding. However, counseling is often brief, standardized, and focused on the transmission of guidelines rather than meaningful engagement with caregivers' lived realities. This limits the ability to address individual concerns, cultural beliefs, and practical barriers.

In addition, inconsistencies in messaging across different healthcare providers can lead to confusion and reduce trust in formal recommendations. When advice appears impractical or disconnected from everyday experiences, caregivers may be less likely to adopt it.

Limitations of Conventional Health Education Approaches

Traditional health education strategies have largely emphasized prescriptive messaging, focusing on what caregivers should do, without adequately addressing why existing practices persist or how recommended behaviors can be realistically adopted. Behavior change is a gradual and complex process that requires trust, dialogue, and adaptation to local contexts.

Programs that fail to engage with the socio-cultural dimensions of feeding practices risk limited effectiveness, even when they are evidence-based. A shift toward more interactive, context-sensitive approaches is therefore essential.

Culture as a Resource: Rethinking the Narrative

It is important to recognize that not all cultural practices are detrimental. Many traditional feeding practices have evolved in response to local environmental conditions and resource availability, and some may align with sound nutritional principles. Framing culture solely as a barrier risks overlooking these positive elements and may alienate communities.

A more effective approach lies in identifying beneficial practices, addressing harmful ones, and integrating scientific recommendations within existing cultural frameworks. Respectful engagement with local knowledge systems can enhance the acceptability and sustainability of interventions.

Way Forward: Toward Context-Sensitive Interventions

Addressing suboptimal weaning practices in rural India requires a shift from generalized, information-driven strategies to approaches that are contextually grounded and culturally responsive. Interventions must move beyond the assumption that knowledge alone can drive behavior change and instead engage with the realities that shape caregiving practices.

A family-centered approach is essential. Given the influential role of elder family members, particularly grandmothers, interventions should extend beyond mothers to include key decision-makers within the household. Engaging these stakeholders can help align traditional knowledge with evidence-based recommendations and reduce intra-household conflicts in caregiving practices.

Equally important is the need for context-sensitive counseling. Frontline healthcare providers should be equipped not only with technical knowledge but also with the skills to deliver culturally competent and empathetic communication. Tailoring advice to local food availability, economic constraints, and household dynamics can enhance the feasibility and acceptance of recommended practices.

Strengthening the capacity of frontline health workers is a critical component of this shift. Training programs should emphasize interpersonal communication, behavioral insights, and cultural awareness, enabling providers to move from directive messaging toward meaningful dialogue with caregivers.

Community-based strategies also offer significant potential. Peer support groups, participatory learning approaches, and sustained engagement within communities can foster trust and facilitate gradual, durable behavior change. Unlike one-time informational sessions, these approaches acknowledge that behavioral transformation is a continuous process shaped by social reinforcement and lived experience.

Conclusion

The persistence of suboptimal weaning practices in rural India underscores the limitations of interventions that rely predominantly on information dissemination. Infant feeding behaviors are shaped by a complex interplay of cultural beliefs, family dynamics, socioeconomic constraints, and health system interactions, all of which influence how knowledge is translated into practice.

Bridging the gap between knowledge and practice requires a paradigm shift, from viewing caregivers as passive recipients of information to recognizing them as active decision-makers operating within specific social and cultural contexts. Integrating scientific recommendations with local realities, rather than attempting to replace them, is key to achieving meaningful and sustainable improvements.

Ultimately, improving infant weaning practices is not solely a question of increasing awareness, but of understanding and engaging with the environments in which caregiving decisions are made. Only through such context-sensitive approaches can lasting improvements in infant nutrition and health outcomes be realized.

References

1. Jha PK, Shukla S, Acharya S. Trends and practices of weaning in infants across India: A comprehensive review. J Family Med Prim Care. 2024 Jul;13(7):2568-2575. doi: 10.4103/jfmpc.jfmpc_1744_23
2. WHO Guideline for complementary feeding of infants and young children 6–23 months of age [Internet]. Geneva: World Health Organization; 2023. 3, Recommendations. Available from: https://www.ncbi.nlm.nih.gov/books/NBK596423/
3. Hanifa IN, Limavady A, Fewtrell M. Exploring factors influencing complementary feeding practices in Asia from parents' and caregivers' perspectives: A qualitative systematic review. Nutr Diet. 2026 Feb;83(1):109-124. doi: 10.1111/1747-0080.70031
4. Sharma M, Gaidhane A, Choudhari SG. A Review of Infant and Young Child Feeding Practices and Their Challenges in India. Cureus. 2024 Aug 9;16(8):e66499. doi: 10.7759/cureus.66499
5. Prakash S, Bindu SA. Infant feeding practices and the associated factors in a rural community in Thiruvananthapuram District, Kerala. J Family Med Prim Care. 2026 Jan;15(1):348-356. doi: 10.4103/jfmpc.jfmpc_609_25
6. Nandagire WH, Atuhaire C, Egeineh AT, Nkfusai CN, Tsoka-Gwegweni JM, Cumber SN. Exploring cultural beliefs and practices associated with weaning of children aged 0-12 months by mothers attending services at Maternal Child Health Clinic Kalisizo Hospital, Uganda. Pan Afr Med J. 2019 Sep 24;34:47. doi: 10.11604/pamj.2019.34.47.16940
7. Lin JL, Lipstein EA, Wittenberg E, Tay D, Lundstrom R, Lundstrom GL, Sediqzadah S, Wright DR. Intergenerational Decision Making: The Role of Family Relationships in Medical Decision Making. MDM Policy Pract. 2021 Oct 29;6(2):23814683211039468. doi: 10.1177/23814683211039468
8. Fikadu K, Takele A, Tesfaye B, Abebo ZH. Intra-household joint decision making on child feeding and associated social determinants in rural districts of South Ethiopia: a multi-site concurrent mixed method study. Front Public Health. 2024 Dec 12;12:1381068. doi: 10.3389/fpubh.2024.1381068
9. Scarpa G, Berrang-Ford L, Twesigomwe S, Kakwangire P, Galazoula M, Zavaleta-Cortijo C, Patterson K, Namanya DB, Lwasa S, Nowembabazi E, Kesande C; IHACC Research Team; Cade JE. Socio-economic and environmental factors affecting breastfeeding and complementary feeding practices among Batwa and Bakiga communities in south-western Uganda. PLOS Glob Public Health. 2022 Mar 9;2(3):e0000144. doi: 10.1371/journal.pgph.0000144

This article is published under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0)